

Asymptotical small- x behaviour of the spin structure functions g_1 and g_2 .

B.I.Ermolaev*

Ioffe Physico-Technical Institute, St.Petersburg 194021, Russia

S.I.Troyan†

Petersburg Nuclear Physics Institute, Gatchina 188350, Russia

Abstract

We show that for small x and in the double logarithmic approximation (DLA) g_2 can be expressed through derivative of g_1 with respect to logarithm of the QCD coupling. Therefore the small- x behavior of the both structure functions is similar. The analytical expression for flavor nonsinglet structure function g_1 is presented and its asymptotical behaviour is calculated.

In framework of the conventional parton model the process of DIS can be considered as a superposition of two independent phases: hadron fragmentation in constituent partons and further DIS of a lepton on one of these partons. At asymptotically high energies, $s \rightarrow \infty$, the behaviour of the cross section is governed by the second phase.

The spin-dependent part of the hadronic tensor for the second phase with the initial parton being a quark, $W_{[\mu\nu]}$, can be written in terms of longitudinal and transversal projections of the quark spin, relative to the plane of the incoming 4-vectors p (quark) and q (virtual photon), in a following way:

$$W_{[\mu\nu]} = i\epsilon_{\mu\nu\alpha\beta} \frac{mq_\alpha}{pq} \left(g_{\parallel} \mathcal{S}_\beta^{\parallel} + g_{\perp} \mathcal{S}_\beta^{\perp} \right) \quad (1)$$

$$g_{\parallel} = g_1, \quad g_{\perp} = g_1 + g_2$$

where m and $\mathcal{S}$ are mass and spin of the quark, and g_1 , g_2 are functions of $x = Q^2/2pq$ and $Q^2 = -q^2 > 0$.

In the Born approximation

$$g_{\perp} = g_{\parallel} = \frac{e_q^2}{2} \delta(1-x), \quad (2)$$

*Supported in part by the Grant INTAS-RFBR-95-311

†Supported in part by the Grant R2G 300 from International Science Foundation

with e_q being the electric charge of the quark. This means that in the Born approximation $g_2 = 0$.

In higher QCD orders at $\ln(1/x) \gg \ln(Q^2/\mu^2)$ all contributions of order of $(\alpha_s(\ln x)^2)^k$ must be taken into full account.

g_1 was calculated in [1, 2] in the DLA analytically for flavour nonsinglet (NS) case and numerically for singlet (S). Here we present the analytical expression for g_1^S and the new result [3] that the structure function g_2 can be obtained through differentiation in QCD coupling α_s the analytical expression for g_1 .

The nonsinglet quark ladder diagram with n gluon rungs in Feynman gauge gives the following contribution to the DLA for antisymmetric structure function

$$W_{\mu\nu}^{(n)} = \prod_{i=1}^n \left[\frac{\alpha_s C_F}{4\pi} \int_x^{\beta_{i-1}} \frac{d\beta_i}{\beta_i} \int_0^{2\pi} \frac{d\phi_i}{2\pi} \int_{\kappa_i}^{s\beta_i} \frac{dk_{i\perp}^2}{[k_i^2]^2} \right] \beta_n \frac{e_q^2}{2s} \delta(\beta_n - x - \frac{\vec{k}_{n\perp}^2}{s}) T_{\mu\nu}^{(n)}, \quad (3)$$

with

$$\kappa_i = \mu^2 + \frac{\beta_i}{\beta_{i-1}} \vec{k}_{i-1\perp}^2$$

where $k_i = \alpha_i q + \beta_i p + k_i^\perp$ is 4-momentum of the quark in a cell i , and

$$T_{[\mu\nu]}^{(n)} = i\varepsilon_{\mu\nu\lambda\sigma} q^\lambda \frac{1}{2} \text{Tr} \left\{ (\hat{p} + m)(1 - \gamma_5 \hat{S}) \gamma_5 \Gamma^\sigma \right\}, \quad (4)$$

where

$$\Gamma^\sigma = \left(\prod_{i=1}^n [-g^{\lambda_i \lambda_i'}] \right) \gamma_{\lambda_1'} \hat{k}_1 \dots \gamma_{\lambda_n'} \hat{k}_n \gamma^\sigma \hat{k}_n \gamma_{\lambda_n} \dots \hat{k}_1 \gamma_{\lambda_1}. \quad (5)$$

The key point for understanding the structure of the matrix Γ_σ is that

$$\hat{k} \gamma^\sigma \hat{k} = -k^2 f^{\sigma\sigma'}(k) \gamma_{\sigma'}, \quad f^{\sigma\sigma'}(k) = g^{\sigma\sigma'} - 2 \frac{k^\sigma k^{\sigma'}}{k^2}. \quad (6)$$

This means that Γ_σ is linear in γ^σ and can be expressed in terms of convolution of n similar f -tensors:

$$\Gamma^\sigma = \left(\prod_{i=1}^n [-2k_i^2] \right) \mathcal{E}_\tau^\sigma \gamma^\tau, \quad \mathcal{E}_{\sigma\tau} = \left(\prod_{i=1}^n \otimes f(k_i) \right)_{\sigma\tau}. \quad (7)$$

Let us present the tensor product $\mathcal{E}$ of Eq.(7) as follows:

$$\mathcal{E}_{\sigma\rho}^{(n)} = g_{\sigma\rho} - \sum_{i=1}^n 2 \frac{(k_i)_\sigma (k_i)_\rho}{k_i^2} + \sum_{i>j}^n 2 \frac{(k_i)_\sigma 2(k_i k_j)(k_j)_\rho}{k_i^2 k_j^2} + \dots \quad (8)$$

As soon as in DLA $k_i^{\perp 2} \simeq k_i^2$, the azimuthal integration of the first two terms of the tensor $\mathcal{E}$ results in

$$E_{\sigma\tau}^{(n)} = \int_0^{2\pi} \frac{d\phi_1}{2\pi} \dots \int_0^{2\pi} \frac{d\phi_n}{2\pi} \mathcal{E}_{\sigma\tau}^{(n)} \simeq g_{\sigma\tau} - n g_{\sigma\tau}^\perp \quad (9)$$

where n is the number of quark cells in the ladder graph.

The analysis in [3] shows that the next terms of the expansion Eq.(8) after integration over azimuthal angles in DLA contribute nothing.

Using the simple relation

$$\frac{\partial \alpha_s^n}{\partial \ln \alpha_s} = n \alpha_s^n$$

one can rewrite the antisymmetric hadronic tensor Eq.(3) as

$$W_{[\mu\nu]}^{(n)} = i \varepsilon_{\mu\nu\lambda\sigma} \frac{m q^\lambda}{(pq)} \left\{ M^{(n)} \mathcal{S}^\sigma - \frac{\partial}{\partial \ln \alpha_s} M^{(n)} \mathcal{S}_\perp^\sigma \right\} \quad (10)$$

with

$$M^{(n)} = \left(\frac{\alpha_s C_F}{2\pi} \right)^n \prod_{i=1}^n \left[\int_x^{\beta_{i-1}} \frac{d\beta_i}{\beta_i} \int_{\kappa_i}^{\beta_i s} \frac{dk_{i\perp}^2}{k_{i\perp}^2} \right] \beta_n \frac{1}{2} e_q^2 \delta(\beta_n - x - \frac{k_{n\perp}^2}{s}). \quad (11)$$

Summing up over all ladders thus leads to a very simple relation between the contributions to DIS structure functions g_1^{NS} and g_2^{NS} :

$$g_2^{NS} = - \frac{\partial g_1^{NS}}{\partial \ln \alpha_s} \quad (12)$$

Eq.(12) is valid only for ladder Feynman graphs. To understand the effect of nonladder graphs on Eq.(12), let us add a nonladder gluon to a ladder graph. Nonladder gluon must be soft enough not to destroy the DL pattern of a ladder graph. Unlike a skeleton ladder gluon a nonladder gluon invokes an additional DL integration only if no any $k_\perp^2$ factor appears in the integrand's numerator. Such factor would cancel the softest virtuality propagator (of those invoked by a nonladder gluon) in the denominator. Thus adding a nonladder gluon does not change the relation between contributions of the $\mathcal{S}^\perp$ and $\mathcal{S}^\parallel$ -structures.

Therefore we would use Eq.(12) if one could separate out the contribution of ladder skeleton gluons to g_1^{NS} and differentiate it w.r.t. α_s . Fortunately one can use the expression for g_1^{NS} obtained in the work [1] and tag the ladder and nonladder gluon contributions through definition of separate QCD couplings α_L and α_{NL} respectively, $\alpha_L = \alpha_{NL} = \alpha_s$. [3]

Then Eq.(12) can be presented in the form

$$g_2^{NS} = - \frac{\partial g_1^{NS}}{\partial \ln \alpha_L} \Big|_{\alpha_L = \alpha_{NL} = \alpha_s} \quad (13)$$

that predicts the same asymptotics of g_1^{NS} and g_2^{NS} at $x \rightarrow 0$:

$$g_2^{NS} \sim g_1^{NS} \sim \left(\frac{1}{x} \right)^a \left(\frac{Q^2}{\mu^2} \right)^{\frac{a}{2}}, \quad a \simeq \sqrt{\frac{2\alpha_s C_F}{\pi}} \left(1 + \frac{1}{2N_c^2} \right). \quad (14)$$

We have considered above the nonsinglet contribution to g_2 . Obviously, one should expect the singlet contribution, i.e. insertion of gluon ladders, to be dominating over the nonsinglet contribution [2]. One could repeat the analysis presented above for the case of gluon ladder contributions, the crucial points of the analysis as well as general features of the result remain the same. The only difference stems from different contributions of a gluon rung to the numerator's tensor structure Eq.(4) for each case: for a quark ladder - Eq.(6) and for a gluon ladder -

$$h^{\sigma\sigma'}(k) = g^{\sigma\sigma'} - \frac{k^\sigma k^{\sigma'}}{k^2}. \quad (15)$$

Thus we obtain the following generalization of the expression (13) for the total structure function g_2^S :

$$g_2^S = - \frac{\partial g_1^S}{\partial \ln \alpha_L} \Big|_{\alpha_L = \alpha_{NL} = \alpha_s} - \frac{1}{2} \frac{\partial g_1^S}{\partial \ln \tilde{\alpha}_L} \Big|_{\tilde{\alpha}_L = \alpha_L = \alpha_{NL} = \alpha_s}. \quad (16)$$

where we differ the QCD coupling $\tilde{\alpha}_L$ corresponding to gluon radiation in a gluon cell from the QCD coupling α_L corresponding to gluon radiation in a quark cell of a ladder graph.

The analytical expression¹ for g_1^S at small x is

$$g_1^S = \frac{\sum e_q^2}{4} \int \frac{d\omega}{2\pi i} \left(\frac{1}{x} \right)^\omega \frac{\omega \left(2\omega + r_1 + r_2 + \frac{b_{(-)}}{h} (r_1 - r_2) \right) \exp(\Omega y)}{\omega^2 + \omega(r_1 + r_2) + r_1 r_2} \quad (17)$$

where

$$\begin{aligned} r_1 &= \sqrt{\omega^2 - 2b_{(+)} + 2h}, & r_2 &= \sqrt{\omega^2 - 2b_{(+)} - 2h}, \\ h &= \sqrt{b_{(-)}^2 + 4b_1 b_2}, & \Omega &= \frac{\omega - r_2}{2}, & y &= \ln \frac{Q^2}{\mu^2}, \end{aligned}$$

and

$$b_{(\pm)} = \frac{\alpha_L}{2\pi} C_F \pm \frac{\tilde{\alpha}_L}{2\pi} 4N_c + \left(\frac{\alpha_{NL}}{2\pi} \right)^2 \left(\frac{2C_F}{N_c} \mp 8N_c^2 \right) \frac{1}{\omega^2}, \quad (18)$$

$$b_1 = \frac{\alpha_L}{2\pi} 2C_F - \left(\frac{\alpha_{NL}}{2\pi} \right)^2 \frac{4C_F N_c}{\omega^2}, \quad b_2 = -\frac{\tilde{\alpha}_L}{2\pi} + \left(\frac{\alpha_{NL}}{2\pi} \right)^2 \frac{2N_c}{\omega^2} \quad (19)$$

where $1/\omega^2$ -terms correspond to nonladder gluon contributions so that color octet amplitudes are taken in the Born approximation. This significantly simplifies formulae but does not noticeably change the small- x asymptotics of g_1^S .

Thus asymptotics of g_2^S and g_1^S at $x \rightarrow 0$ is

$$g_2^S \sim g_1^S \sim \left(\frac{1}{x} \right)^{\omega_0} \left(\frac{Q^2}{\mu^2} \right)^{\frac{\omega_0}{2}} \quad (20)$$

¹In the previous work [2] the small- x asymptotics of g_1^S was calculated numerically.

where ω_0 is the rightmost singularity (branching point) of the integrand in Eq.(17):

$$\omega_0 = 2b_{(+)}(\omega_0) + 2h(\omega_0). \quad (21)$$

This is an algebraic equation of the fourth order. Solution of this equation gives $\omega_0 \simeq 3.5\sqrt{\alpha_s/2\pi}$. If one neglects quark contributions to the ladder then $\omega_0 \simeq 3.7\sqrt{\alpha_s/2\pi}$. The small difference between these results shows that similar to Pomeron gluons give the dominating contribution to asymptotics of g_2^S and g_1^S .

Let us recall that the above results were obtained in the DLA. Worthwhile to notice that taking single log effects into account, though does not change the power-like character of the Eqs.(14,20), changes the values of the powers. For g_1^{NS} the new power is

$$a' = a \left[\left(1 + \frac{a^2}{16} \right)^{\frac{1}{2}} + \frac{a}{4} \right]. \quad (22)$$

For example, if one takes $\alpha_s = 0.18$, the Eq.(22) gives: $a \simeq 0.4$, $a' \simeq 0.44$.

References

- [1] J.Bartels, B.I.Ermolaev and M.G.Ryskin, Z.Phys.C 70 (1996) 273.
- [2] J.Bartels, B.I.Ermolaev and M.G.Ryskin, Z.Phys.C 72 (1996) 627.
- [3] B.I.Ermolaev and S.I.Troyan, Preprint PNPI TH-14-1997 2157;
hep-ph/9703384